



# RapiSol<sup>®</sup>

## 3-2-1

Mn

Cu

Zn

**EDTA Technology**  
**Micro Nutrient**  
WETTABLE GRANULE (WG)



9 341200 000232

**20Kg**  
(4x5kg)

**A 100% Manganese, Zinc and Copper EDTA technology formulation to prevent and correct deficiencies in the crops specified on the label.**

**Manganese (Mn) as EDTA: 7.8% w/w**  
**Zinc (Zn) as EDTA: 4.2% w/w**  
**Copper (Cu) as EDTA: 1.5% w/w**

**DIRECTIONS FOR USE:**

- Do not apply successive foliar applications within 14 days, unless otherwise indicated.
- When applied with other EDTA foliar fertiliser products, the total amount must not exceed the maximum hectare rate for each individual product for a specific crop.
- DO NOT apply EDTA technology formulations to broadacre crops by foliar application greater than 1%, i.e. 1 kg per 100 L.
- Do not mix with acid based products nor acidic water, as crop damage may result.
- When applied to soils, ensure the soil pH is within the range of 4 to 9.

| CROP  | RATE                       | APPLICATION    | CRITICAL COMMENTS   |
|---|----------------------------|----------------|---|
| Cereals,<br>Grain Legumes,<br>Oilseed crops | 0.25-1 kg/ha<br>1-3 kg/ha  | Foliar<br>Soil | Apply when symptoms appear.<br>Apply with other fertilizers in the sowing operation.  |
| Cotton, Maize,<br>Summer crops              | 0.5-1.5 kg/ha<br>1-3 kg/ha | Foliar<br>Soil | Apply from 4-6 leaf stage of the crop and/or when symptoms appear.<br>Apply with other fertilizers in the sowing operation. |
| Pastures,<br>Lucerne                        | 0.75-1.0<br>kg/ha          | Foliar         | Apply when symptoms appear.   |

**MIXING INSTRUCTIONS:**

Add RapiSol 3-2-1 to water while agitating. Continue agitation until the granules have fully dissolved. Do not pre-mix.

**COMPATIBILITY**

RapiSol 3-2-1 is compatible with foliar fungicides (e.g. propiconazole and tebuconazole formulations), Awaken® ST, NutriSync™ M, NutriSync™ D, Carfentrazone-ethyl, MCPA and 2,4-D based herbicide formulations.

Improved uptake may be achieved with the addition of Citowett®, Flexstic®, Flextend® and Hygro-Stic® adjuvants.

Do not apply with organosilicone type surfactants or other cuticle disruptive adjuvants. Compatible with most commonly used fertilizers and agrochemicals. If unsure, it is recommended to perform a jar test before mixing, followed by a small scale application before widespread use.

**WARNING:** Keep out of reach of children. Harmful if swallowed. If the product comes into contact with skin or eyes, wash immediately with plenty of water. Store in a fresh, dry place and protect from excessive heat. Store the product in its original container. Use the product according to the instructions on the label. If in doubt, please contact your local agronomist. If local experience is unavailable, it is recommended a small scale test application is conducted before widespread use.

**USE ONLY IN CASE OF PROVEN NECESSITY. DO NOT EXCEED THE MAXIMUM DOSAGES.**

**CONDITIONS OF SALE:** AGSPEC warrants that the product within the unopened pack conforms to the chemical description on this label. As the occurrence of unusual environmental conditions and the use of this product, is all beyond the control of AGSPEC, no guarantee, expressed or implied, is made as to the effects of such use or the results obtained, and in no event shall the liability of AGSPEC exceed the price of the product used. The purchaser must assume all risks of storage, use or application of this product. Goods are sold subject to the standard Conditions of Sale of the manufacturer, available on request. Use of the product implies that the purchaser accepts these conditions of sale.

Packed and Distributed in Australia by  
**Agspec Australia Pty Ltd.**  
ABN: 401 095 739 53  
PO. Box 1006, Mt, Gambier, S.A., 5290  
Toll Free- 1800 683 456  
Email- info@agspec.net  
Website- www.agspec.net

