



RapiSol[®]



EDTA Technology
Micro Nutrient
WETTABLE GRANULE (WG)



20kg

A 100% Copper (Cu) EDTA technology wettable granule formulation to prevent and correct copper deficiency in the crops specified in the label.

Copper (Cu) as EDTA: 14 % w/w

DIRECTIONS FOR USE:

- Do not apply to fruit crops by foliar application greater than 0.1% concentration (i.e. 100 gm per 100 L) or to broadacre crops by greater than 1% (i.e. 1 kg per 100 L).
- To avoid defoliation risk in fruit trees, do not exceed the 0.1% concentration, i.e. 100 g/100 L.
- Caution is required when applied during flowering when other products are included, as the outcome from mixed products maybe unknown.
- Do not apply during fruit colour change.
- Do not apply successive foliar applications within 14 days, unless otherwise indicated below.
- When applied with other EDTA foliar fertiliser products, the total amount must not exceed 0.75 kg/ha (or 0.5 kg/ha for broadacre crops, 1.0 kg/ha for grapes and tree crops, or 2 kg/ha for Forestry).
- When applied to soils, ensure the soil pH is within the range of 4 to 9.
- Do not mix with acid based products nor acid water, as crop damage may result.

Crop	Rate	Application	Critical Comments
Cereals, Grain Legumes, Oilseed crops, Cotton, Maize, Summer crops	0.1-0.5 kg/ha	Foliar	Apply when symptoms appear.
	0.75-1 kg/ha	Soil	Apply with other fertilizers in the sowing operation and before tillering.
Pastures, Lucerne and Forage crops	0.5-0.75 kg/ha	Foliar	Apply when symptoms appear.
	0.75-1 kg/ha	Soil	Apply with fertilizers in planting row, or in a separate application before pasture emergence.
Potatoes and other tuber crops	0.5-0.75 kg/ha	Foliar	Apply when symptoms appear. Apply in a minimum of 500 L/ha of water.
Vegetables	15-40 g/100 L	Foliar	Apply when sufficient foliage is present. Apply in a minimum of 500 L/ha of water.
	0.5-1.5 kg/ha	Soil	Apply as a broadcast spray application before sowing, at transplanting or beginning of plant growth.
Grapevines (wine and table)	50-100 g/100 L or 0.5-1.0 kg/ha	Foliar	Apply early in the crop cycle. Dilute application only. Do not exceed maximum per hectare rate. Minimum water rate of 500 L/ha.
	2-5 g/vine	Soil	Usually one application should be sufficient. This rate can be split into smaller applications throughout the season.
Apples, Nut crops, Citrus, Mangoes, Avocados, Pineapples, Olives	50 g/100 L or 0.5-1.0 kg/ha	Foliar	Apply early in the crop cycle. Dilute application only. Do not exceed maximum per hectare rate. Minimum water rate of 500 L/ha.
	2-5 g/tree	Soil	Usually one application should be sufficient. This rate can be split into smaller applications throughout the season.
Strawberries, Other berry crops	0.5-0.75 kg/ha	Foliar	Apply when symptoms appear. Apply in a minimum of 500 L/ha of water.
	2-3 kg/ha	Soil	Apply as a broadcast spray application before sowing, at transplanting or beginning of plant growth.
Forestry (including pines and blue gums)	1-2 kg/ha	Foliar	Apply when symptoms appear. Apply in a minimum of 200 L/ha of water.
Ornamentals	10-40 g/100 L or 0.5-0.75 kg/ha	Foliar	Apply early in the season. Apply in a minimum of 200 L/ha of water.
	3-5 g/bush	Soil	Apply before sowing, at transplanting or beginning of plant growth.

MIXING INSTRUCTIONS

Add RapiSol Cu to water while agitating. Continue agitation until the granules have fully dissolved. Do not pre-mix.

COMPATIBILITY

RapiSol Cu is compatible with foliar fungicides (e.g. propiconazole and tebuconazole formulations), Awaken® ST, NutriSync™ M and NutriSync™ D, Carfentrazone-ethyl, MCPA and 24D based herbicide formulations.

Improved uptake may be achieved with the addition of Citowett®, Flexstic®, Flextend® or Hygro-Stic® adjuvants.

Do not apply with organosilicone type surfactants or other cuticle disruptive adjuvants.

Compatible with most commonly used fertilisers and agrochemicals. If unsure, it is recommended to perform a jar test before mixing, followed by a small scale application before widespread use.

WARNINGS

Keep out of reach of children. Harmful if swallowed. If the product comes into contact with skin or eyes, wash immediately with plenty of water. Store in a fresh, dry place and protect from excessive heat. Store the product in its original container. Use the product according to the instructions on the label. If in doubt, please contact your local agronomist or representative. If local experience is unavailable, it is recommended a small scale test application is conducted before widespread use.

USE ONLY IN CASE OF PROVEN NECESSITY. DO NOT EXCEED THE MAXIMUM DOSAGES.

CONDITIONS OF SALE: AGSPEC warrants that the product within the unopened pack conforms to the chemical description on this label. As the occurrence of unusual environmental conditions and the use of this product, is all beyond the control of AGSPEC, no guarantee, expressed or implied, is made as to the effects of such use or the results obtained, and in no event shall the liability of AGSPEC exceed the price of the product used. The purchaser must assume all risks of storage, use or application of this product. Use of the product implies that the purchaser accepts these conditions of sale.

Packed and Distributed in Australia by

Agspec Australia Pty Ltd.

ABN: 401 095 739 53

PO. Box 1006, Mt. Gambier, S.A., 5290

Toll Free- 1800 683 456

Email- info@agspec.net

www.agspec.net



Citowett®, Flexstic®, Flextend®, Hygro-Stic® and RapiSol® are registered trademarks of Agspec Australia Pty Ltd.

Nutrisync™ is a trademark of Loveland Products, Inc.

Awaken® is a registered trademark of Loveland Products, Inc.

