



# RapiSol<sup>®</sup>

Zn



**EDTA Technology**  
**Micro Nutrient**  
WETTABLE GRANULE (WG)



**20kg**

**A 100% Zinc (Zn) EDTA Technology wettable granule formulation to prevent and correct zinc deficiency in the crops specified in the label.**

**Zinc (Zn) as EDTA: 14 % w/w**

**DIRECTIONS FOR USE:**

- Do not apply to fruit crops by foliar application greater than 0.1% concentration (i.e. 100 gm per 100 L) or to broadacre crops by greater than 1% (i.e. 1 kg per 100 L).
- Caution is required when applied during flowering when other products are included, as the outcome from mixed products maybe unknown.
- Do not apply during fruit colour change.
- Do not use by foliar application on plum or apricot trees.
- Do not apply successive foliar applications within 14 days.
- When applied with other EDTA foliar fertiliser products, the total amount must not exceed 1.0 kg/ha (or 1.5 kg/ha for vegetable crops).
- When applied to soils, ensure the soil pH is within the range of 4 to 9.

Crop	Rate	Application	Critical Comments
Cereals, Grain Legumes, Oilseed crops, Cotton, Maize, Summer crops, Pastures, Lucerne and Forage crops	0.25-1 kg/ha 1-3 kg/ha	Foliar Soil	Apply from early post emergence, and/or when symptoms appear. Apply with other fertilizers in the sowing operation.
Potatoes and other tuber crops	0.5-1.5 kg/ha	Foliar	Apply when symptoms appear. Apply in a minimum of 500 L/ha of water.
Vegetables	0.5-1.5 kg/ha	Foliar	Apply when sufficient foliage is present. Apply in a minimum of 500 L/ha of water.
	2-4 kg/ha	Soil	Apply as a broadcast spray application before sowing, at transplanting or beginning of plant growth.
	2-3 kg/ha	Fertigation	Total dosage split into several applications during the crop cycle. Apply by drip/trickle irrigation or microjet sprinklers to the soil.
	0.3-0.5 kg/1000 L	Hydroponics	Apply as required-1 L of prepared solution per 100 L irrigation water.
Grapevines (wine and table)	33 - 100 g/100 L or 0.5-1.0 kg/ha	Foliar	Apply 2-3 times during the crop cycle. Do not exceed 3x concentration. Use lower rate when concentrate spraying. Do not exceed maximum per hectare rate. Minimum water rate of 250 L /ha.
	2-3 kg/ha	Fertigation	Total dosage split into several applications during the crop cycle. Apply by drip/trickle irrigation or microjet sprinklers to the soil.
Apples, Pears, Nut crops, Citrus, Mangoes, Stonefruit, Avocados, Pineapples, Olives	50 g/100 L or 0.5-1.0 kg/ha	Foliar	Apply 2-3 times during the crop cycle. Dilute application only. Do not exceed maximum per hectare rate. Minimum water rate of 500 L/ha.
	2-3 kg/ha	Fertigation	Total dosage split into several applications during the crop cycle. Apply by drip/trickle irrigation or microjet sprinklers to the soil.
Strawberries, other berry crops	0.5-1.0 kg/ha	Foliar	Apply when sufficient foliage is present. Apply in a minimum of 500 L/ha of water.
	4-8 kg/ha	Soil	Apply as a broadcast spray application before sowing, at transplanting or beginning of plant growth.
	2-3 kg/ha	Fertigation	Total dosage split into several applications during the crop cycle. Apply by drip/trickle irrigation or microjet sprinklers to the soil.
	0.3-0.5 kg/1000 L	Hydroponics	Apply as required-1 L of prepared solution per 100 L irrigation water.
Ornamentals	75-150 g/100 L	Foliar	Apply early in the season.
	4-8 kg/ha	Soil	Apply as a broadcast spray application before sowing, at transplanting or beginning of plant growth.
	2-3 kg/ha	Fertigation	Total dosage split into several applications during the crop cycle. Apply by drip/trickle irrigation or microjet sprinklers to the soil.
	0.3-0.5 kg/1000 L	Hydroponics	Apply as required-1 L of prepared solution per 100 L irrigation water.

**MIXING INSTRUCTIONS**

Add RapiSol Zn to water while agitating. Continue agitation until the granules have fully dissolved. Do not pre-mix.

**COMPATIBILITY**

RapiSol Zn is compatible with foliar fungicides (e.g. propiconazole and tebuconazole formulations), Awaken® ST, NutriSync™ M, NutriSync™ D, Carfentrazone-ethyl, MCPA and 2,4-D based herbicide formulations.

Improved uptake may be achieved with the addition of Citowett®, Flexstic®, Flextend® or Hygro-Stic® adjuvants.

Do not apply with organosilicone type surfactants or other cuticle disruptive adjuvants.

Compatible with most commonly used fertilizers and agrochemicals. If unsure, it is recommended to perform a jar test before mixing, followed by a small scale application before widespread use.

**WARNINGS**

Keep out of reach of children. Harmful if swallowed. If it comes into contact with skin or eyes, wash immediately with plenty of water. Store in a fresh, dry place and protect from excessive heat. Store the product in its original container. Use the product according to the instructions on the label. If in doubt, please contact your local agronomist or representative. If local experience is unavailable, it is recommended a small scale test application is conducted before widespread use.

**USE ONLY IN CASE OF PROVEN NECESSITY. DO NOT EXCEED THE MAXIMUM DOSAGES.**

**CONDITIONS OF SALE:** AGSPEC warrants that the product within the unopened pack conforms to the chemical description on this label. As the occurrence of unusual environmental conditions, and the use of this product, is all beyond the control of AGSPEC, no guarantee, expressed or implied, is made as to the effects of such use or the results obtained, and in no event shall the liability of AGSPEC exceed the price of the product used. The purchaser must assume all risks of storage, use or application of this product. Use of the product implies that the purchaser accepts these conditions of sale.

Packed and Distributed in Australia by  
**Agspec Australia Pty Ltd.**  
 ABN: 401 095 739 53  
 PO. Box 1006, Mt, Gambier, S.A., 5290  
 Toll Free- 1800 683 456  
 Email- info@agspec.net  
 www.agspec.net



Citowett®, Flexstic®, Flextend®, Hygro-Stic® and RapiSol® are registered trademarks of Agspec Australia Pty Ltd.  
 NutriSync™ is a trademark of Loveland Products Inc.  
 Awaken® ST is a registered trademark of Loveland Products Inc.